

EXPERTS FOR GROWTH



Basfoliar® K Premium SL



Foliar Fertiliser
Bio-powered liquid formulae



Basfoliar® K Premium SL

Foliar Fertilizer, Bio-powered Liquid Formula

Basfoliar K Premium SL is a liquid foliar fertilizer with high potassium concentration, boosted with extracts of Seaweed (*Ecklonia maxima*) and Boron.

The combination of elements that make up Basfoliar K Premium SL favors:


- Color, flavour and size of the fruit.
- Formation and mobilization of carbohydrates towards the fruit (sugars).
- Stimulation of photosynthesis.
- Post harvest life of fruits stored in cold.
- Tolerance to cold climates, diseases and resistance to stress due to lack of water.

The formulation of Basfoliar K Premium SL (liquid) and the high quality of its materials prevents risks of stained fruit, even applied the day before harvest.

Composition :

Potassium (K ₂ O)	25% p/v
Ecklonia maxima (Algal extract)	3% p/v
Boron	1% p/v

Electrostatic Charge :



No Problems of Capping in Nozzles

As seen in the graph, Basfoliar K Premium SL presents an excellent behavior in applications with electrostatic machine, maintaining without alteration a adequate electrical charge in the solution and without problems of obstruction in the nozzles during the application.

Recommended doses:

Crop	No. of application	Doses	Time of application
Grapes (Table & wine)	2 to 3	2.5 to 3.0 ml/ Lt.	7-15 days interval during fruit development and maturity stage
Apple	3	2.5 to 3.0 ml/ Lt.	Apply every 15 to 21 days (60, 40 and 20 days before harvest)
Pomegranate	3 to 4	2.5 to 3.0 ml/ Lt.	Apply at least 3 to 4 times every 15 days, interval during fruit development. Last application 10 days before harvest.
Melons/Cucurbits	2 to 3	2.5 to 3.0 ml/ Lt.	Apply, at fruit development and fruit maturity stages
Citrus	2 to 3	2.5 to 3.0 ml/ Lt.	Apply every 15 days from the fruit set.
Strawberries	2 to 3	2.0 - 2.5 ml/Lt.	Apply from the first fruit set, every 10 to 12 days
Tomato	4 to 6	2.5 to 3.0 ml/ Lt.	Apply from the set of the 2nd to the 5th cluster every 5 to 7 days.
Garlic and Onions	2	2.5 to 3.0 ml/ Lt.	Apply 30 and 15 days before harvest
Vegetables & Potatoes	2 to 3	2.5 to 3.0 ml/ Lt.	Apply from 1/3 of crop development on the row, every 10 to 15 days.
Flowers/Ornamental plants	2 to 3	2.0 ml/Lt.	Apply at 7 to 10 days interval after flowering

Precautions:

Basfoliar K Premium SL is an environmentally compatible and biodegradable product.

Basfoliar K Premium SL is considered non-toxic for plants and animals. For safe handling, it is necessary to abide by the precautions for the use of plant protection products with which Basfoliar K Premium SL has to be mixed.

